

SCIENCE

A WEEKLY RECORD OF SCIENTIFIC PROGRESS.

ILLUSTRATED.

Entered in the Office of the Librarian of Congress, at Washington, D. C.

No. 46—Vol. II.

May 14, 1881.

Price 10 Cents.

THE "ACME" MICROSCOPES.



"Acme" No. 2.

Monocular, . . . \$100
Binocular, . . . 141


"Acme," No. 3.

Monocular, . . . \$45
Binocular, . . . 70

"Acme" No. 4.

The favorite Laboratory
Stand, . . . \$25
With 1 inch and $\frac{1}{4}$ inch
objectives, . . . 45

Special Terms made with
Colleges.



"ACME" No. 4.



"Acme" Lithological.

A new Stand, after the pattern of the No. 3, for the study of Rocks; convertible in an instant into an ordinary monocular.

All our Stands

have same size of tubing and sub-stage fitting, so that eye-pieces and accessories that fit one Stand can be used with all.

"Congress" Turn-Table
Patented, \$6.50.

Gem Turn-Table, \$2.50.
Do. Do. with improved
centering adjustment, \$3.25



HOLMAN'S COMPRESSOR, THE LATEST AND THE BEST, \$4. MECHANICAL FINGER, \$5
JOHN W. SIDLE & CO., Lancaster, Pa.

Send for Catalogue.

D. APPLETON & CO.,

HAVE RECENTLY PUBLISHED:

General Physiology of Muscles and Nerves.

By B. I. ROSENTHAL, Professor of Physiology in the University of Erlangen. With Seventy-five woodcuts. ("International Scientific Series.") 12mo, cloth. Price \$1.50.

"This attempt at a connected account of the General Physiology of Muscles and Nerves is, as far as I know, the first of its kind. The general data for this branch of science has been gained only within the past thirty years."—*Extract from Preface.*

Popular Lectures on Scientific Subjects.

By H. HELMHOLTZ, Professor of Physics in the University of Berlin. Second series. 12mo, cloth. Price, \$1.50.

CONTENTS.

- I. Gustav Magnus. In Memoriam.
- II. On the Origin and Significance of Geometrical Axioms.
- III. On the Relation of Optics to Painting—Form, Shade, Color, Harmony of Color.
- IV. On the Origin of the Planetary System.
- V. On Thought in Medicine.
- VI. On Academic Freedom in German Universities.

The favor with which the first series of Professor Helmholtz's lectures were received justifies, if a justification is needed, the publication of the present volume.

Animal Life,

AS AFFECTED BY THE NATURAL CONDITIONS OF EXISTENCE. By CARL SEMPER, Professor of the University of Würzburg. With Maps and One Hundred Woodcuts. "International Scientific Series." 12mo, cloth. Price \$2.00.

"It appears to me that of all the properties of the animal organism, Variability is that which may first and most easily be traced by exact investigation to its efficient causes; and as it is beyond a doubt the subject around which, at the present moment, the strife of opinions is most violent, it is that which will be most likely to repay the trouble of closer research. I have endeavored to facilitate this task, so far as in me lies." *From the Preface.*

"This is in many respects one of the most interesting contributions to zoological literature which has appeared for some time."—*Nature.*

SIGHT.

AN EXPOSITION OF THE PRINCIPLES OF MONOCULAR AND BINOCULAR VISION. By JOSEPH LE CONTE, LL.D., author of "Elements of Geology," etc. ("International Scientific Series.") With Illustrations. 12mo, cloth. Price, \$1.50.

"In writing this treatise I have tried to make a book that would be intelligible and interesting to the thoughtful, general reader, and at the same time profitable to even the most advanced specialist in this department."—*From Preface.*

The Atomic Theory.

By AD. WURTZ, Member of the French Institute. Translated by E. CLEMINSHAW, M. A., Assistant Master at Sherborne School. Number XXIX. of "The International Scientific Series." 12mo, cloth, price, \$1.50.

The Power of Movement in Plants

By CHARLES DARWIN, LL. D., F. R. S., assisted by FRANCIS DARWIN. With Illustrations. 12mo, 592 pages, cloth, price, \$2.00.

"Mr. Darwin's latest study of plant-life shows no abatement of his power of work or his habits of fresh and original observation. We have learned to expect from him at intervals, never much prolonged, the results of special research in some by-path or other subordinated to the main course of the biological system associated with his name; and it has been an unflinching source of interest to see the central ideas of the evolution and the continuity of life developed in detail through a series of special treatises, each we might say exhaustive of the materials available for its subject."—*Saturday Review.*

For sale by all booksellers; or sent by mail, to any part of the United States, post-paid, on receipt of price.

D. APPLETON & CO.,

Publishers,

1, 3 and 5 Bond Street,

NEW YORK.

CLUB RATES.

	Subscription.	With SCIENCE.		Subscription.	With SCIENCE.
Analyst	\$2.00	\$5.50	Iron	7.50	9.34
Appleton's Monthly	3.00	6.26	Journal of the Telegraph	2.00	5.50
AMERICAN NATURALIST	4.00	7.20	London Lancet	5.00	7.50
AMERICAN JOURNAL OF SCIENCE	6.00	8.50	Magazine of American History	5.00	8.00
American Journal of Microscopy	1.00	4.70	MINING AND SCIENTIFIC PRESS	4.00	6.40
American Machinist	3.00	6.70	Nature	6.00	9.15
American Monthly Microscopical Journal	1.00	4.80	N. Y. Medical Journal	4.00	7.00
American Manufacturer and Iron World	4.25	7.00	New England Journal of Education	3.00	6.40
Boston Journal of Chemistry	1.00	4.75	North American Review	5.00	7.75
Chicago Medical Review	2.00	5.50	Operator	1.00	4.80
Chicago Field	4.00	7.00	POPULAR SCIENCE MONTHLY	5.00	8.00
Druggist	1.60	5.25	Sanitary Engineer	3.00	6.50
Druggists' Circular	1.50	5.25	Practical American	1.50	5.00
Educational Monthly (Barnes)	1.50	5.20	Pacific Rural Press	4.00	7.00
Engineering News	3.00	6.25	Scribner's Monthly	4.00	7.20
Engineering and Mining Journal	4.00	7.00	Science Observer	.50	4.40
Education	4.00	7.00	SCIENTIFIC AMERICAN	3.20	6.50
Engineering Magazine	5.00	8.25	" " SUPPLEMENT	5.00	8.00
Harper's Magazine	4.00	7.00	" " WITH SUPPLEMENT	7.00	9.60
Harper's Weekly	4.00	7.20	Southern Medical Record	2.00	5.40
Harper's Bazar	4.00	7.20	Student	1.00	4.75
Harper's Young People	1.50	5.20	St. Nicholas	3.00	6.40
HUMBOLDT LIBRARY	3.00	6.25	Young Scientist	.50	4.30
International Review	5.00	8.00			

SCIENCE :

A WEEKLY RECORD OF SCIENTIFIC
PROGRESS.

JOHN MICHELS, Editor.

PUBLISHED AT
TRIBUNE BUILDING, NEW YORK.
P. O. Box 8888.

SATURDAY, MAY 14, 1881.

THE alleged discovery of a new motive force for driving engines, patented by Professor Gamgee, of Washington, is already condemned on theoretical grounds, both in this country and in Europe.

The principle involved is not a new one, and, so far, all previous attempts in the same direction have ended in failure. In this case, Chief Engineer Isherwood, of the United States Navy, gives an endorsement to Professor Gamgee's scheme, which has caused some eminent physicists to give an attention to it which perhaps it hardly deserved.

From what we can gather, we understand that Professor Gamgee proposes to work his engine with ammonia, taking advantage of the fact that in a liquified state it boils at -37.3° Fahr., and that at 60° Fahr. it exerts a pressure of seven atmospheres—or, say, 100 lbs. to the square inch.

Authorities differ on this subject, but so far as liquid ammonia is concerned, it is stated as follows: "That at atmospheric pressure, and a temperature of 62° Fahr., 1 lb. of the gas occupies about 23 cubic feet, while 1 lb. of liquid ammonia would occupy only 36 cubic inches."

According to Mr. Isherwood, the "zero-motor" is an apparatus in which liquid ammonia can be vaporized under considerable pressure by means of the heat in water, or in the external atmosphere, and the gas so obtained is used to propel a piston through a cylinder—the gas being employed with the greatest measure of expansion found possible.

At this point the difficulty is presented of returning the ammonia to the boiler. Professor Gamgee offers no explanation, but claims to be able to accomplish it by some method he has invented. He asserts that in its expansion the liberated gas is refrigerated and diminished in bulk, and becomes partially liquified at the end of the stroke of the piston, when it is exhausted and returned from whence it came.

Against this, Professor Simon Newcomb and some English writers assert, that in the absence of

demonstration to the contrary, it will absorb as much power to convert the ammonia gas into the liquid form as the latter will give out when vaporized.

In the "zero-motor" Professor Gamgee professes to have an engine capable of exerting great power, and without the necessity of using any fuel, and indirectly the claim is made of solving successfully the problem of perpetual motion.

Apart from some fundamental errors which underlie the scheme, many theoretical difficulties could be suggested, but as a practical test of "the discovery" will probably be made, further discussion may profitably be postponed until the result of the trial is known.

It will no doubt be a genuine surprise to all students of nature to learn that a German scientist has found fossil plants and animal forms in most of the meteorites (chondrites) which he has examined for the purpose.

DR. OTTO HAHN, who has taken a prominent part in the discussion on the "*Eozoön canadense*" has, in the usual way, prepared sections of many of these bodies. These he has had photographed and thereby attained a result which is independent of the microscopist's vision. Dr. Hahn claims that they show many forms of plants and animals in a fossil state contained in their mass, of which the highest forms are crinoids, corals and allied species. He has placed this collection of sections in the hands of Dr. Weiland of Tübingen, (formerly of Philadelphia) for thorough classification.

We regret that we are unable to endorse this interesting discovery. Professor Whitfield, superintendent of the fossils and minerals in the American Museum of Natural History, has seen Dr. Hahn's drawings and was unable to verify the presence of the organic forms referred to. He attributed Dr. Hahn's error to a too sanguine temperament, and an "imagination which bodies forth the form of things unknown."

WE are indebted to our Washington correspondent for a brief mention of an interesting paper by Dr. George M. Sternberg, on "A Fatal Form of Septicæmia in the Rabbit, produced by Subcutaneous Injection of Human Saliva."

Dr. Sternberg recently published a translation of Dr. Antoine Magnin's work on Bacteria, and has had considerable experience in making investigations on septic organisms. He now asserts that the human saliva carries with it a deadly poison, which will kill a rabbit in forty-eight hours; other animals also appear to be influenced more or less by the same cause, while still others—the dog, for instance—resist the

poison. Some salivas are more fatal than others—that of Dr. Sternberg being especially virulent. It will be noticed in our report that Dr. Sternberg attributes the poisonous element to the presence of Micrococci—having found this form of Bacteria both in the saliva employed and in the poisoned blood of the victims.

These facts may be considered in conjunction with experiments by M. Pasteur in the same direction.

SCIENTIFIC SOCIETIES IN WASHINGTON.

THE BIOLOGICAL SOCIETY.—At the last two meetings the Society has listened to four papers: A Fatal Form of *Septicæmia* in the Rabbit produced by Subcutaneous Injection of Human Saliva, by Dr. George M. Sternberg; On the Mortality of Marine Animals in the Gulf of Mexico, by Mr. Ernest Ingersoll; A Statistical View of the Flora of the District of Columbia, by Professor Lester F. Ward; and Notes on Scale Insects, by Professor J. H. Comstock. All of these papers were of the highest scientific value, prepared by specialists in connection with their own immediate investigations. Dr. Sternberg has been making experiments for the past two years under the patronage of the National Board of Health, concerning the causes and development of epidemic diseases. In the course of his labors he has made careful observations with reference to inoculation, and in the paper referred to above gave the Society the benefit of his experiments on saliva injected under the skin of the rabbit. As an elaborate report will appear in the proceedings of the Board of Health, it will be necessary to state only the conclusions arrived at, which are as follows: The rabbits impregnated died invariably in less than 48 hours. Other animals which did not succumb were afflicted with sores. Dogs resist the poison, guinea pigs yield less readily than rabbits, fowls escape entirely. Some salivas are more fatal than others; that of Dr. Sternberg is especially virulent. The presence of Micrococci in the saliva, in the blood of the poisoned animals, and in that of animals infected with this poisoned blood led the author to the conviction that the evil effect was owing to these minute bacteria.

Mr. Ernest Ingersoll, who has been studying the waters of the Gulf of Mexico in the interest of the U. S. Fish Commission, reported that in certain years there occurred a great mortality among the marine animals. In the years 1844, 1854 and 1878 such disasters had been noticed, but the one most injurious in its consequences was in the year 1880. Oysters, clams, fish, and even sponges, were involved in the universal ruin. The beaches were so thickly covered with the dead bodies that the inhabitants were driven from their homes. Various attempts were made to account for the phenomenon, but with indifferent success.

Professor Ward is preparing a work to be entitled "A Catalogue of the Flora of the District of Columbia." It will include all the phænogamous plants and the vascular cryptogams. The number of species enumerated is 1233, distributed among 526 genera, as follows:

Polypetalous	genera, 173	Species, 354
Gamopetalous	" 169	" 388
Monochlamydeous	" 47	" 122
Monocotyledonous	" 120	" 321
Coniferæ	" 4	" 7
Vascular Cryptogams	" 13	" 41
	526	1233

Professor Ward then proceeded to give the census of these species with reference to the orders, to the position of the district north and south, and east and west, as well as in comparison with local floras which have been described with sufficient accuracy.

Professor Comstock's paper on the *Coccida*, or scale insects, was a very entertaining treatment of a very dry

subject. The group under discussion is usually regarded as the most uninteresting of all the animal kingdom as well as the most anomalous. It is true that the lac of commerce and that of Arizona is the product of these insects, but the most of them are worthless or pernicious. They infest greenhouse plants and most of our useful fruit and timber trees. A specimen was exhibited which had been taken from Europe to Los Angeles, Cal., and back to Washington, upon a lemon, and at the end of its nine-thousand-mile trip was as lively as ever. The method of hatching, of the deposit of the meal, or lac, and of moulting in the male and female, were described and illustrated with drawings and cabinet specimens. The method of classifying these animals into species has been a very uncertain one. Even the later used characteristic, namely, the series of pores or openings on the penultimate ring not being always invariable. Professor Comstock has found the fringe on the last segment of the abdomen to be the most constant specific characteristic. An interesting point in the paper was a discovery made by Mrs. Comstock, that the poisers behind the wings are furnished with a hooklike process which fits into a groove on the back of the wing and helps to sustain it in flight.

THE ANTHROPOLOGICAL SOCIETY.—The entire session of the Society at its last meeting was occupied with the reading and discussion of a paper read by the Rev. Clay McAuley upon the Seminole Indians still remaining in Florida. Of this once formidable but now humbled tribe there remain in the vicinity of Lake Okeechobee 208 individuals, 37 families, 22 camps, and 5 settlements. There are no half-breeds among them, the occurrence of such a birth would probably subject the author to torture or death. They are healthy, have an abundance of food, and are probably increasing. The men are tall, well proportioned, erect, lithe, and graceful. The women are shapely, agreeable, vigorous, and many of them handsome. Dr. McAuley singled out three, whom he characterized as the stately, the beautiful, and the handsome among all the Indians whom he had visited. A very minute description was given of the dress, ornament, customs, and language of these people. A full report will appear in the publication of Major J. W. Powell's Bureau of Ethnology.

THE ULTRA-GASEOUS OR RADIANT STATE OF MATTER.*

BY PROF. H. S. CARHART.

The announcement by Mr. William Crookes, F. R. S., some six years ago, that he had produced mechanical motion by the direct impact of waves of light created a profound impression in the scientific world. But when it was found that the Radiometer, which was supposed to exhibit this new action of light, carried a system of blackened vanes, delicately balanced in a very high vacuum, it impressed most physicists as being an interesting form of heat engine receiving its supply of heat by absorption of radiant energy.

Mr. Crookes subsequently adopted the same view, which was the only tenable one, and investigated the subject in a long series of exceedingly skilful and ingenious experiments. His researches on the Radiometer formed the introduction to a more extended series of investigations into the movement of the residual gas of very high vacua, under the influence of heat and the negative discharge of electricity. These investigations carry us to the very farthest boundary of matter thus far attained, and furnish an ocular demonstration of some of those molecular movements that have heretofore been merely imagined. In fact the phenomena observed are such that Mr. Crookes has felt himself justified in announcing a fourth, or ultra-gaseous state of matter. Such an an-

*Lecture delivered before the New York Electrical Society on May 5, 1881.

nouncement startles us by its boldness, and encounters the opposition of our conservatism. I shall have the pleasure of reproducing a part of Mr. Crooke's experiments before you with tubes made after his models, and recently imported; and at the close of the exhibition I hope you will be able to draw your own conclusions.

Let us first prepare the way from our study of the subject by a clear presentation to our minds of the exact differences between the solid, the liquid, and the gaseous states. We shall then be able the better to judge whether this new claimant is sufficiently differentiated from ordinary gases to warrant its being set aside by itself.

I. A solid is composed of distinct molecules separated from one another by spaces which are large compared with the diameter of molecules themselves. These molecules, which are the indivisible units so far as all *physical* changes are concerned, are held rigidly in a fixed relation to one another by cohesion, a force which acts only across these invisible inter-molecular spaces, and the intensity of which depends upon the chemical constitution of the body. The force of cohesion is counterbalanced by the motion of the individual molecules, the energy of this motion constituting the heat of the body. As the temperature rises, the amplitude of vibration increases, and the force of cohesion is diminished by an increase of distance between the molecules. When the temperature falls the reverse process takes place.

Molecules of matter, in the solid state, retain fixity of position about their centres of oscillation; and any attempt to change the relation of these centres, by distorting the body, calls into action resistances opposing the change of form.

II. When the temperature rises so high that the molecules lose their fixity of position, and are released from the rigid thralldom of cohesion, so that they can apparently roll round one another without changing the distances of their centres, then the solid becomes liquid. As the force of cohesion differs with each kind of matter, so the temperatures of liquefaction also differ. In liquids the cohesion is much diminished but not entirely overcome by the energy of molecular oscillations, which we call heat. A liquid is characterized by taking the form of the containing vessel, by its surface becoming horizontal, and by its equal transmission of applied pressure in all directions. The state of liquidity is due to intermolecular motions of much greater amplitude than those peculiar to the solid state.

III. The temperature still rising, the molecules are finally released entirely from the force of cohesion, and move about in every conceivable direction with enormous and constantly varying velocities. The mean distance between them is sufficiently great to render cohesion inoperative; but they come into constant collisions with one another, now colliding, and now rebounding, changing their direction of motion and perhaps their velocity with every impact. On account of this rapid molecular motion and the absence of cohesion in gases, they tend to expand indefinitely; and they transmit pressure by multitudinous collisions among the molecules, and exert pressure on the walls of the containing vessel by innumerable impacts against its interior. Place a small, closed balloon, partly filled with air, under the receiver of an airpump and exhaust. The balloon distends more and more as the exhaustion proceeds, because of this molecular bombardment against its inner surface, the external counteracting pressure being withdrawn. In much the same way an impenetrable sheet of metal might be kept suspended horizontally in the air by bullets fired vertically against its under surface. The dynamical principles concerned in the two cases are identical. When a gas is heated, the velocity of the individual molecules increases, and they exert greater pressure on the walls of the containing vessel by their impact, or by expansion secure a longer free path for themselves between successive collisions with one another. This mean free path, or mean

length of path, is known to be about $\frac{1}{1000000}$ of an inch at the average or normal pressure of the air, while the mean distance between the molecules is about one seven-millionth of an inch, according to the late Prof. Clerk Maxwell. Accordingly a cubic inch of air at normal density must contain the cube of 7,000,000 or 343 quintillions of molecules. Each of these molecules, moreover has, perhaps, millions of encounters every second, the number varying with the kind of gas considered. The state of gaseity is therefore pre-eminently characterized by innumerable molecular collisions, from which proceed all the properties constituting the ordinary gaseous state of matter; and this state continues so long as the motions of the molecules are in every possible direction, and the collisions almost infinite in number. Under such conditions the mean length of path of the molecules, or the distance traversed between two successive collisions, is extremely small compared with the dimensions of the containing vessel, only $\frac{1}{1000000}$ of an inch.

Such are the distinctions existing between the solid, the liquid, and the gaseous states, distinctions depending entirely upon molecular aggregation. In the snow or frost crystal, for instance, the molecules of water are grouped together in a definite structural relation; the application of heat destroys the structure, the molecules becoming more nearly individualized by a certain increase of independence or freedom of motion, and the solid becomes a liquid; a still further rise of temperature increases this molecular independence to such an extent that their only physical relationship depends upon innumerable collisions with one another. If now, by any means, molecular independence can be rendered so nearly complete that the hits may be disregarded in comparison with the misses, then the free path of the molecule between the hits may become comparable with the dimensions of the containing vessel, the properties which constitute gaseity are reduced to a minimum, and matter is exalted to an ultra-gaseous state. The residual gas is then in that peculiar state when it has ceased to have the power of adjusting its own pressure; and consequently the phenomenon of diverting the molecules by suitable means into any paths at desire will be possible. Such is the ultimate result of gaseous expansion. As the rarefaction proceeds the number of molecules in a given space becomes smaller and smaller, and the mean length of path greater and greater, the molecular collisions diminishing as the free path increases.

At an exhaustion of a millionth of an atmosphere the number of molecules in any given space is reduced to one millionth of the number at the ordinary pressure; each molecule has a million times more room to move about in, and its mean length of path becomes about four inches (1,000,000 times $\frac{1}{1000000}$ in.) If now the rarefaction be carried to another millionth, the space appropriated to each molecule is then increased a million million times, but the distance between the molecules becomes only one seven-hundredth of an inch, and a cubic inch contain seven then not less than 340 millions of molecules, according to Prof. Clerk Maxwell. This rarefaction is about 50,000 times further than the best mercurial air-pump can attain. A gas transmits pressure instantaneously in all directions by innumerable molecular collisions; an *ultra gas*, or "radiant matter," has lost the power of adjusting its pressure because of the infrequency of collisions among the molecules.

If then a bulb about four inches in diameter is exhausted to a millionth of an atmosphere, the residual gas in it will have lost the power of rapid adjustment of pressure to equality at every point. Hence any extraneous force, like heat or electricity, may "infuse order into the apparently disorderly jostling of the molecules in every direction by coercing them into a methodical rectilinear movement. * * * * And according to the extent to which this onward movement has replaced the irregular motions, which constitute the very essence of the

gaseous condition, to that extent have the molecules assumed the condition of radiant matter."

Let us now see to what extent Mr. Crookes has reached these conditions in his Radiometer and radiant matter tubes.

In the Radiometer the blackened vanes become heated by absorbing radiant energy (both heat and light), and they project the contiguous gaseous molecules from their surfaces by communicating to them their molecular motion, much as a vibrating drum-head would project into air grains of sand strewn on it. An increase of pressure is thus produced between the heated lamp-black and the gaseous residue; but this pressure is not transmitted throughout the whole bulb because of the relative infrequency of collisions among the molecules at this degree of exhaustion. Hence the molecules are projected forward against the cooler bulb in right lines, and the blackened vanes retreat because the repulsion between them and the gaseous residue is mutual. [A Radiometer projected on the screen by an oxyhydrogen lantern; also, a diagram showing the paths of the projected molecules].

Mr. Crookes investigated the Radiometer with an ingenuity and a variety of detail that left nothing to be desired. Moreover, his visualization of the molecular motions taking place in this little instrument prepared him for a further research into the motion of the residual gas in Geissler tubes. The dark space around the negative pole, which broadens as the exhaustion proceeds, and the brilliant stratification displayed by many Geissler tubes, were hints in the line of his special studies. I cannot forbear, at this point, to call your attention to the vibration of the air in a sounding pipe, made visible by very light, precipitated silica powder. The resemblance between the beautiful segmentation of the pipe by the thin, vertical planes of silica powder, and the stratification of a Geissler tube, is most marked and suggestive. [A finely stratified Geissler tube shown; and a glass pipe, containing silica powder, and vibrated by a whistle projected on the screen]. As the silica planes limit the free swing of the air particles in the pipe, so the bright portions of the vacuum tube show when the residual gases of contiguous segments encounter each other—the vibration being set up by the passage of electricity. And it may be observed here that as the transmission of sound is the onward transference of energy by motion, so the passage of electricity is the transference of energy from one point to another by means of another form of motion.

The exhaustion which answers best for a Geissler tube is comparatively low. I have frequently obtained beautiful stratification in a tube six feet long, exhausted by a good air-pump. Mr. Crookes carried the exhaustion much further and obtained entirely new results. The first noticeable feature was the gradual broadening of the dark space surrounding the negative pole as the exhaustion advanced [dark space tube, exhibited a transverse sheet of aluminum in the middle constituting the negative pole]. This dark space is regarded as the free path of the molecules at this degree of exhaustion. It increases as the exhaustion proceeds, and contracts when the exhaustion diminishes. The molecules of air are projected normally from the negative pole, and the illumination at the boundary of the dark space is due to the collision between the gas projected from the negative pole and the more slowly moving molecules advancing toward it. Here, then the lines of molecular pressure, caused by the excitement of the negative pole, are illuminated by the induction spark.

With still higher vacua the free path becomes equal to the dimensions of the containing vessel, and the projected molecules impinge directly against its walls. The molecules stream from the negative pole with enormous velocities and dart across the tube with comparatively few collisions. Their motion is then arrested by the solid matter of the tube. A noteworthy property of this radiant

matter then appears. Luminosity is produced by the impact of the projected molecules against solid matter. Phosphorescence, as this luminosity is called, is thus excited, its color depending upon the kind of matter receiving the impact. [Three phosphorescent tubes shown; also a bulb partly filled with phosphorescing material.] English glass phosphoresces a light blue, uranium glass rather dark green, and soft German glass, light apple green. Rubies always shine with a deep red hue whatever the color of the gem itself. The artificial rubies made in Paris show no variation from the real stones in the color of their phosphorescent light. This phosphorescence takes place better at an exhaustion of about a millionth of an atmosphere than at any other. A tube designed to show the dependence of the phosphorescence upon exhaustion, is made by connecting to the main tube a small supplementary tube containing caustic potash, which holds captive a certain amount of aqueous vapor [the tube exhibited]. Turning on the coil, the tube now shows green phosphorescence, but upon heating the potash tube with a small lamp, aqueous vapor is released, the phosphorescence gradually disappears, and is replaced by the stratified discharge of a Geissler tube. Withdrawing the lamp, the vapor is re-absorbed by the caustic potash, the fine stratification widens out slowly and finally retreats towards the potash bulb, while a wave of green light sweeps from the negative pole, driving the last pale stratification into the potash tube. Radiant matter moves in straight lines and absolutely refuses to turn a corner. This peculiarity is shown by a V-shaped tube. [The tube exhibited.] The flood of green light proceeds from the negative pole at the top only as far as the bottom, refusing to turn the angle toward the positive in the other branch. Reversing the current, the illuminated branch follows the negative pole. A striking contrast is thus presented between a Geissler and a Crookes tube; but this contrast is brought out more clearly still by two bulbs exactly alike, except in the degree of their exhaustion. Both are fitted with one negative terminal and three positive ones. With the low vacuum tube the stratification follows the direction of the positive pole, changing its path as that pole is changed. Changing now to the high vacuum or Crookes tube, containing only about a millionth of an atmosphere, and turning on the coil the only light to be seen is the green phosphorescence of the glass. The negative pole is a very shallow cup, and the projected radiant matter crosses its focus and then strikes on the opposite side of the bulb where the impact excites strong phosphorescence. Changing the positive pole produces no change whatever in the path of the projected molecules or the luminosity of the glass. The positive pole exercises no influence whatever upon the direction of discharge of radiant matter from the negative pole. The residual gas is here exalted to the fourth or radiant State, and its free path, under the impulsion of the negative discharge, is entirely across the bulb. Moreover, not only is luminosity induced, but the glass bulb rapidly heats where it receives the cannonade of these invisible balls.

Another peculiarity of radiant matter, depending upon its projection in right lines, is that when intercepted by solid matter it casts a shadow. This large pear-shaped bulb has the negative pole at the small end. A cross cut out of sheet aluminum is placed across the bulb so as to intercept a part of the gaseous molecules streaming from the negative pole. Connecting with the induction coil, the dark shadow of the cross is seen plainly projected on the large end of the bulb. Here the projected molecules pass by the aluminum cross and bombard the walls of the bulb, producing the usual phosphorescence. Nothing could show more distinctly than the projection of matter from the negative pole in straight lines.

This bombarding causes the surface of the glass to lose its sensitiveness to intense phosphorescence. The cross is hinged so as to turn down with a slight shake.

Again turning on the coil the rays from the negative pole fall uninterruptedly on the large end of the bulb; the dark cross is now replaced by one more intensely green than the adjacent portions of the glass. The fresh parts of the glass are more susceptible to luminous excitation than the areas previously bombarded. This stenciled image of the black cross, Mr. Crookes tells us, is so persistent that it remained in one case after the glass had been heated hot, so that the end of the bulb was bent in and then blown out again. After re-exhaustion the bright green cross came out plainly in the more intense phosphorescence produced by the electrically projected residual gas. Thus the molecules hammer away upon the glass with sufficient energy to produce a permanent impression.

Since the molecules of the residual gas are driven violently from the negative pole, there should be a recoil of the pole from the molecules. That such is the case is shown by this electrical radiometer. The connections are such that the aluminum vanes constitute the negative pole. The vanes are not blackened but are covered on one side with mica, a non-conductor. The uncovered sides constitute the radiating or projecting surfaces. Turning on the induction current the vanes are repelled and the system rotates rapidly. The phosphorescent spots on the glass, produced by the impact of the molecules projected from the vanes, rotate as the vanes do, giving us a visible image of the process going on in the heat radiometer. In this second form the negative pole consists of a ring of platinum wire. Above this is a fly, composed of mica vanes inclined like the fans of a wind-mill. Turning on the current the matter projected from the wire strikes against the sloping vanes and sets them in motion. But this is not all. I now connect a battery directly with the terminals of the platinum ring, and the passage of the current heats the ring red hot. Directly the fly begins to turn more rapidly than before. Radiant matter is thus projected by heat as well as electricity, and the ordinary heat radiometer is propelled by the recoil of projected radiant matter like the electrical radiometer.

The action of magnetism on this stream of electrified particles is, indeed, curious. When a straight Geissler tube of low vacuum is employed it gives a narrow line of violet light joining the two poles. [This tube placed over the poles of an electro-magnet.] On passing the battery current through the magnet underneath, the violet line is repelled or attracted according to the direction of the currents; but it recovers itself after passing the magnet and proceeds to the other pole. Not so with a tube of high vacuum. This tube contains a mica screen, covered with material which phosphoresces under molecular impact. The radiant matter from the negative pole passes through a narrow opening and impinges upon the screen along its entire length. You observe how the phosphorescence marks the path of the projected molecules. Bringing a strong magnet down over the stream or actuating the electro-magnet, and the luminous path curves toward the side of the tube like the path of a projectile; but in this case the stream does not recover its original direction after deflection, as in the case of the Geissler tube. This same tube is fitted to determine another question of much interest connected with these wonderful phenomena. It is provided with two negative projecting surfaces, and the stream of molecules from each of these may be made to trace its path on the mica screen in a luminous line. Observe the position of this line first from one pole and then the other. What now will be the result if streams issue from both poles at once? If they constitute two currents of electricity in the same direction then we know that they will attract each other; but if they consist of a train of similarly electrified molecules, then they must repel each other. Put to the test of experiment we see plainly that the stream lines diverge by mutual repulsion.

We have seen that mechanical action is produced by

the recoil from the radiant matter. It is equally true that strong mechanical effects may be produced by the impact of these swiftly moving molecules. This tube is ingeniously constructed with a pair of glass rails running from end to end. Along them rolls freely the axle of a small wheel with broad vanes as paddles. The poles are so situated that the radiating molecules may strike against the vanes. Turning on the induced current, the stream of swiftly moving molecules strikes against the vanes on one side of the wheel and sets it running along the rails. Reversing the current, the wheel stops and returns on its track. [Tube projected on a screen by oxy-hydrogen light.] Another tube has been devised to show both magnetic deflection and mechanical action on the screen. The negative pole is a large, shallow cup. A mica screen intercepts the converging streams of radiant matter.

Behind the screen is placed an easily-revolving mica wheel provided with vanes and making a sort of paddle wheel. So arranged the molecular rays from the pole are cut off from the wheel, and no movement is produced. Placing a magnet over the tube the rays are deflected so as to pass above the screen and the wheel begins to revolve. Reversing the magnetic poles, the deflection is in the other direction, and the wheel now rotates like an undershot water-wheel. Even at this high vacuum of about a millionth of an atmosphere, enough matter remains in the tube to produce a sort of molecular wind under the impulsion of the negative discharge. Radiant matter produces heat as well as mechanical motion when its motion is arrested. The negative pole in this tube is a cup or concave projector. The radiant matter comes to a focus near the cup and then passes on toward the other end of the tube. But on bringing a magnet near it the stream of radiant matter is deflected so as to strike the side of the tube, producing a green, phosphorescent spot. The nearer the magnet approaches the greater the deflection of the stream and the smaller the spot. Covering the side of the tube with wax and placing it in front of the lantern, a dark image of the tube is projected because of the opaque wax. But with the coil in action the approach of the magnet deflects the radiant stream, the tube is heated by the impact of the molecules, the wax melts and becomes transparent, and the light from the lantern passes through.

The heat generated by the arrest of radiant matter is further shown by a bulb of special device. It has a negative pole of aluminum in the form of a concave projector. In its focus is fixed a piece of platinum foil. Turning on the coil, the projected molecules impinge against the platinum at last raise it to a red heat. Indeed Mr. Crookes has actually melted iridio-platinum in such a focus. We are accustomed to see powerful heating effects produced by currents of electricity traversing rather poor conductors; but here red hot platinum glows with the invisible cannonade of innumerable molecules of the air we breathe.

These effects take place indifferently with hydrogen, carbonic acid gas, and air at this high vacuum. The only difference appears to be that the phosphorescence begins at different pressures with the different gases. The results obtained depend, therefore, not on chemical properties, but on the physical, molecular condition of the gaseous residues. These phenomena are so entirely different from those obtained at ordinary degrees of exhaustion, that Mr. Crookes appears to me to be justified in considering that he has at last verified Faraday's early hypothesis of matter in a radiant state. Mr. Crookes has well said that, "In studying this fourth state of matter we seem at length to have within our grasp and obedient to our control the little indivisible particles which, with good warrant, are supposed to constitute the physical basis of the universe. We have seen that in some of its properties radiant matter is as material as this table, while in other properties it almost assumes the character of radiant energy. We have actually touched the

borderland where matter and force seem to merge into one another, the shadowy realm between the known and the unknown, which for me has always had peculiar temptations. I venture to think that the greatest scientific problems of the future will find their solution in this borderland, and even beyond; here, it seems to me, lie ultimate realities, subtle, far-reaching, wonderful."

GREAT ASTRONOMICAL TELESCOPE.

The greatest refracting telescope in the world—Lord Rosse's is a reflecting telescope—has been constructed for the Vienna Observatory by Mr. Howard Grubb, at his celebrated manufactory of astronomical instruments, at Rathmines, near Dublin. We give an illustration, from a sepia drawing of it by Mr. G. Browning, and abridge the following account of it from a description received at this office.

The idea of crowning the observatory at Vienna with a refracting telescope of surpassing power was conceived by the Austro-Hungarian Government about five years ago. Such a building was worthy of the best instrument that could be constructed. Every visitor to the Austrian capital must be struck by it, standing upon a site of between fourteen and fifteen acres at a height of 200 feet above the city, and extending 330 feet in length and 240 feet in width. Desiring to possess the finest telescope which could be procured, the Government commissioned Dr. Edward Weiss, now Director-General of the Observatory at Vienna, to visit all the principal observatories and workshops in the world. He recommended that the task should be confided to Mr. Grubb, of Dublin, who was ordered to construct a refracting telescope of at least 26 inches aperture. A commission was appointed by the Austro-Hungarian Government to superintend the work. It was composed of the following gentlemen: The Earl of Crawford and Balcarres, Dr. Higgins, the Earl of Rosse, Professor Stokes, of Cambridge, Professor Ball, Astronomer Royal for Ireland, Dr. Stoney, Secretary of the Queen's University in Ireland, many years connected with Lord Rosse's observatory, Dr. E. Reynolds, professor of chemistry, Trinity College, Dublin, and Mr. Walsh, Austro-Hungarian Consul in Dublin. On the 16th ult. the Commissioners reported their unanimous approval of the finished instrument.

The general form of the telescope is that known as Grubb's modified Gramme, and is similar to the well-known standard equatorial which he constructed for the Earl of Crawford and Balcarres, Dr. Huggins, Oxford University, Berlin, Cork, and other places. It possesses all the modern improvements and special arrangements of an ingenious character, which are rendered desirable by its great size. The length of the tube is 33 feet 6 inches, and the aperture is 27 inches. The tube is entirely of steel, $3\frac{1}{2}$ feet in diameter in the centre, and tapering to each end. The entire moving parts, including the tube, polar, and declination axis, counterpoise and various adjustments weigh between six and seven tons; yet the whole apparatus is under such control that one person can move it about and manipulate it with the utmost ease. The mechanism is remarkable for its solidity and strength, as well as for its exquisite delicacy.

In order to render the motion of such ponderous instruments sufficiently easy, the makers are generally obliged to reduce the diameter of the axes, particularly that known as the declination axis, to an extent that makes one almost alarmed for their safety, to say nothing of their stability. Mr. Grubb, however, has mastered the difficulties of the position by a peculiar and most interesting system of equipoise, by which he is enabled to make his axes so large and solid as to ensure stability and give perfect confidence without sacrificing the ease of motion. The application of antifricition apparatus to the polar axis has been already successfully effected, and was a simple prob-

lem, but Mr. Grubb has the exclusive merit of applying it to the declination axis, which is a task of great and complicated difficulty, demanding the highest scientific skill.


Another remarkable feature in the work is the ingenious arrangement by which the circle can be read with the utmost ease and certainty. It is usually a very troublesome operation with large telescopes to read the circle, and when the circles are about 20 feet or more from the ground the labor and delay which it involves are very formidable. In Mr. Grubb's instrument, the circles are carefully and accurately divided on a band of gold, and by a system of reflectors, at once beautifully simple and ingenious, the observer can without stirring from his chair read all the circles of the instruments through one little reader telescope attached to the side of the main telescope tube.

The setting of the telescope is massive and graceful. The frame on which it rests down to the ground level is of cast iron, and there are chambers of considerable size at the base. In the lower one, which is entered by a door at the end, is a clock for driving the instrument in order to follow the paths of the heavenly bodies. The castings of which the frame is formed are of about ten tons weight, and are of simple but not inelegant design. The clock-work is controlled by Mr. Grubb's novel frictional governor, and is also furnished with his new electric control apparatus. There are two right ascension circles, each 2 feet in diameter, one read from the eye end of the telescope and the other from the ground floor. The declination circle is 5 feet in diameter, and is read from the eye end of the telescope. All the circles are divided on an alloy of half pure gold and half pure silver, which is found to be very white and not liable to corrode or tarnish.

The material for the object glasses was procured from M. Feil, of Paris. The protracted delay in procuring this material for the work was a subject of great anxiety to Mr. Grubb, and occasioned heavy additional outlay on his part. In October, 1879, however, discs were obtained, which in working gave good promise, and in December last he was able to report the work finished—his part of it being, in fact, accomplished in less than half the time stipulated by the agreement with the Austro-Hungarian Government. His task was practically trebled by the difficulty experienced in obtaining pure discs. The success of his undertaking is regarded with great satisfaction and with national pride. He has supplied equipments to most of the modern observatories, but this telescope is his greatest achievement.

THE *Athenæum* prints an interesting extract from a letter from Prof. Draper, giving a description of the progress he has made in photographing the nebula in Orion:—"I have succeeded," he says, "in taking stars in it of the 14.1, 14.2, and 14.7 magnitudes of Pogson's scale. Prof. Pickering has made a series of measures on these magnitudes especially for me at the Harvard College Observatory. You will perceive that we have photographed stars which approach the *minimum visible* of my 11 in. telescope, and we may, therefore hope shortly to photograph stars actually too faint to be seen with the eye in the same instrument. The nebula, which was exposed 104 minutes, extends over an area of about 15' in diameter, though, as it becomes fainter toward the exterior parts, it is difficult to determine its precise limits." This is a great advance; no star of less than the ninth and a half magnitude has hitherto been photographed.

A CURIOUS magnetic property of the meteoric iron of Santa Cattarina (Brazil), has been lately observed by Professor Lawrence Smith. Small detached fragments, not weighing more than 0.1 to 0.2 gr., were very weakly affected by a magnet; but on being flattened on a piece of steel, with a steel hammer, they become very sensitive to it. By heating red-hot, the particles were made to be still more easily attracted than by flattening. The meteoric iron in question contains 66 iron, 34 nickel.



THE NEW TELESCOPE, VIENNA OBSERVATORY.

THE POLARIZATION OF SOUND.

AN EXAMINATION INTO THE NATURE OF VIBRATIONS IN EXTENDED MEDIA.*

BY S. W. ROBINSON,

Professor of Physics and Mechanics, Ohio State University.

The phenomena of polarization of light have heretofore presupposed transversal vibrations of particles of the luminiferous ether. Such vibrations have not only been supposed transversal when polarized, but primarily, or when in their primitive condition. It is proposed now to show that no necessity exists for considering the vibrations in light as transversal in front of a polarizer; and furthermore to show that in all probability luminous vibrations are primarily longitudinal.

It is well known that light can be radiated, reflected, refracted, diffracted, diffused, can be made to interfere and can be polarized. All these effects are known to be common to sound, except the last, and it is for the sole purpose of explaining this in light that the convenient theory of transversal vibration has been set up by physicists, for the single case of luminous vibration. It is to be noticed that transversal vibrations are not to be assumed impossible when sufficient cause exists. It is simply assumed that the cause is insufficient when a material particle is made to vibrate from the action of a disturbance at a remote single centre transmitted to the particle considered; the centre, the transmission, and the particle considered, being supposed as belonging to a homogeneous medium of indefinite extent. As regards the nature of the vibratory movements of particles of luminiferous ether may we not justly ask that, if we can go through such a range of density as from platinum to hydrogen without a change in the nature of the vibrations where, as we rise in the scale of ethereal tenuity, shall longitudinal end and transversal begin? Why should the luminiferous ether, now considered as a substance, have a peculiar form of vibration? If ether undulations can be polarized, why not undulations generally? These questions are not answered by the highest authorities. The short of it all seems to be that if polarized light had never been discovered probably the device of transversal vibrations never would have been set up. Indeed, the eminent author M. J. Jamin, says in his three volume work on Physics at the outset, in his lesson on polarization, and subsequent to the treatment of interference, diffraction and other phenomena: "What has been said previously of the movement of luminous waves is absolutely independent of the directions of the vibration." This is good authority for limiting the *transversal* theory to polarization, authority with which doubtless all physicists will agree on this point. It is, therefore, only necessary to polarize sound to place all the known effects of luminous waves in common with sound waves, or to make the theory of longitudinal vibrations universal.

Assent to the above notions will be the more readily given after noticing the consideration that, in polarized light it is not necessary to suppose the vibrations transversal till after passing the polarizer, and that the latter probably imparts an effect equivalent to a lateral impulse, as due to its one-sided action upon the ray transmitted, thus giving cause for vibration which are more or less transversal; the same being true for undulations in air, water, iron, platinum, hydrogen, luminiferous ether, etc., without exception.

But independent of all questions of polarization we find powerful evidence of the unity of system, for vibrations in all possible media; evidence which, in one case at least, is employed as the basis of a rigorous mathematical demonstration of the impossibility of existence of perpetual motion. This latter named evidence is the famous

principle of Helmholtz regarding the action of natural forces among mutually interacting material points, viz.: that the forces must be central forces and functions of the distance, and hence motions of remote particles can only be longitudinal with reference to the centre of force. This principle, considered aside from luminiferous ether, will be universally accepted as truth. But what is the criterion for making an exception of any homogenous substantial medium; even luminiferous ether? Indeed, if any criterion exists for such an exception, it consists simply in a desired convenient means for explaining polarization: a theory of transverse vibrations, which, though convenient and probable beyond the polarizer, has thoughtlessly and without need, been extended to the front of the polarizer, and to undulations in the primitive conditions, where, as shown above, no necessity exists for transversal vibrations. That the necessity for transversal vibrations in primitive rays of light is entirely wanting, let it be granted for the sake of an argument that the source of light, such as the sun or a gas burner, is capable of exciting vibrations in the adjacent medium, which are in all possible directions immediately at the radiant. At a considerable distance from the radiant, the effect upon a single particle will be the resultant of action of all the particles immediately surrounding the radiant, and transmitted by and through the intervening medium; such a resultant impulse can hardly be admitted to be otherwise than longitudinal. Or again, a particle near the radiant imparts an impulse to the adjacent particle. If the passive particle is in direct line with the active one, the impulse received will be in direct line also, and the direction of motion of the two particles coincides. The similar direct action of a second particle upon a third will also be in the same line, and so on indefinitely. At a distance of several hundred or thousand times the diameter of the radiant the line of vibration indicated will be almost perfectly longitudinal irrespective of where the first particle considered is situated about the radiant. That is to say, a particle at the surface of a radiant vibrating in a line of direction tangential to it or transversal, may be considered as transmitting its vibration from particle to particle in a direct line, and hence to the best advantage; and still, at a distance this direct line becomes nearly a line of longitudinal vibration for a remote particle.

Thus all lines of direction of vibration will pass through the radiant or be tangent to it, so that in sunlight the rays, and the directions of vibration, will all lie within the visual angle of the sun, or within about half a degree of arc, and hence almost perfectly longitudinal.

These considerations all confirm the principle of Helmholtz. And that remarkable principle, together with all considerations presented above, and all which can possibly result from a careful study of the subject of transmitted impulses, go to confirm a universal law of longitudinal vibrations for primitive rays in all possible substantial media, and to antagonize the notion of transversal vibration.

Hence, if light can be polarized, why not undulations generally. The writer, after much study of the subject, became convinced of the possibility of this about eight years ago, and six years ago apparatus was made for putting the matter to an experimental test. Want of time prevented, and further study determined a modification of the apparatus which was made over two years ago. This apparatus was successful in verifying all my preconceived notions in the matter, but owing to extended study and matured views of the principles involved, the experiments simply confirmed, without developing new theories or unanticipated facts. I propose now to describe the apparatus and give the results.

The means adopted for polarizing the undulations is the same as that for polarizing light by reflection. It is well known that when sound passes from one medium into another whose velocity of sound differs, the sound is

*Copyright 1881, by S. W. Robinson.

refracted. Recent investigations of Henry, Tyndall and others have indicated that when sound encounters a change of density of medium, as when passing from clear atmosphere into a wall of fog, there is a reflection of sound. Altogether there seems no doubt but sound acts like light in these respects, that is, on meeting a change of refractive power, it is both reflected and refracted, as light is at the surface of water or of glass. The reflected light is found to be always more or less polarized, perfectly so for the so-called polarizing angle. This polarizing angle of incidence is such that, as discovered by Brewster for light, the reflected and refracted components of a single ray, as they strike away from the point of incidence are at right angles. As reflected light is polarized, reflected sound was supposed to be also.




FIG. 1.

Applying the laws of Fresnel and Brewster—1st, that the index of refraction is equal to the ratio of the velocities of the waves in the media; and 2d, that complete polarization is obtained for the particular case of right angled reflected and refracted component rays, we are guided to the proper conditions. We conclude that any two substances having different velocities of propagation of waves may be selected. For instance, two gases, like hydrogen and air, any two liquids, any two solids, a solid and a gas, or, generally, any two media whatever. Considerations of convenience would indicate air and illuminating gas, and these were chosen for the present purpose. The velocities of propagation in air and coal gas being as 1125 and 1420, the index of refraction, according to the first law above, is $n = 1.26$. The second law gives for the polarizing angle of incidence, $\tan i = n = 1.26$, or $i = 51\frac{3}{4}^\circ$, the rays or waves being in the gas. To realize this incidence upon a surface of separation between the gas and air, the coal gas was placed in L-shaped tubes, *AB*, Fig. 1, having a portion cut away at the angle, as shown at *CD*. The branches of the L make equal angles of $51\frac{3}{4}^\circ$ with the normal to *CD*. A delicate membrane was gummed to the tube covering the opening at *CD*, also shown at *F*, the object of which was to retain the gas and maintain a polarizing surface, *CD*. The arrow at *A* indicates a ray which is incident at *E*, and is then in part refracted outward at *E* in a direction perpendicular to the reflected component *EB*. Each tube was about one inch in diameter and three inches long. A number of these were made of tin, each with one end slightly larger than the other, so that they could be joined up, stove-pipe fashion, to any desired extent. Being cylindrical, the plane of one L-piece could be placed at any angle with the plane of the preceding one, according to the desired polarizing test.




FIG. 2.

Fig. 2 shows the manner of joining the tubes, giving

the effect of nine polarizing surfaces, like nine plates of glass in light arranged at the polarizing angle. The nine plates of glass can be used in two parts—one part, 4 for instance, serving as a polarizer and the remaining 5 as analyzer. The ends at *A* and *B* were capped with membranes and the whole filled with illuminating gas. Thus *AC* may serve as a polarizer and *CB*, or *CE*, or *CD* as analyzer. When arranged as in *ACB* or *ACE* all conspire to the same effect of polarization; but when arranged as in *ACD*, the plane of all the L-pieces in *CD* being at right angles to that of those in *AC*, the effect of one part antagonizes that due to the other, and to a maximum degree as regards the angle. Partial effects may be obtained with intermediate angles between 0° and 90° . Again, we observe that the L-pieces of Fig. 2 may be alternately crossed, so that no two contiguous ones will be parallel. It is believed that this arrangement will give the greatest possible antagonistic effect; also while all Ls are in one plane it is not necessary that they be arranged in a zig-zag line, like *AC* and *CE*, but may be indiscriminately connected in that plane. The few experiments made with the above-named arrangements gave very marked results. Of course it need not be confined to nine or any particular number of the L-pieces.




FIG. 3.

It was found, however, wanting in convenience. The apparatus finally adopted is that shown in Fig. 3. A different number of L-pieces were used at different times. The portion *AB* is the polarizer and *BC* the analyzer. The joint at *B* was kept tight with beeswax; the ends at *A* and *C* were capped square with the same membrane material as were the angles of the Ls, giving, when charged with illuminating gas, a continuous zig-zag column from *A* to *C*. The L-pieces of the polarizer enter half Ls at *A* and *B*, the latter having a common axis and resting in bearings at *A* and *B* in the standards, as shown. The object of this is to enable the experimenter to turn the polarizer readily from cross to parallel, etc. This convenient arrangement of the polarizer is due to my assistant, Mr. Wright. Although applied to the polarizer, it is evidently equally applicable to the analyzer instead. The half L angles were not covered with membranes, but left solid, with gradual inside curvature. Membranes might have been applied here with partial polarizing effect. The half L solid angles are supposed to have detracted in a measure from the percentages of polarization obtained; but this sacrifice is more than compensated for by the greater convenience and constancy of conditions obtained. If this arrangement gives decisive results, of course, more perfect apparatus would. The illuminating gas was admitted by a nipple and rubber hose at *C*, the same flowing the length of tubes and issuing in a small jet at *H*; my assistant kept this ignited, and used the flame length as a pressure indicator, and it served admirably.

The first trials were by blowing an organ pipe in front of the membrane *A*, to agitate the gas column.

A small mirror was attached to the membrane *C*, reflecting a pencil of light upon a screen. The deportment of the image indicated complex and inadmissible vibratory movements of gas column, and besides quantitative indication was found preferable to qualitative; thereupon the quantitative impulse and indicator pendulums were adopted, as shown at *E* and *F* respectively, Fig. 3. An ivory ball, $\frac{1}{4}$ inch in diameter, suspended by a thread of 8 inches length, was used at *E*, and so placed that when at rest the ball would just touch the membrane at *A*. The impulse was imparted by bringing the ball back against the stop, shown by means of a spatula held in the hand, and then allowing it to swing free against the membrane, each time with a definite, predetermined arc. So much of the impulse as reaches *C* knocks the pendulum *F* through a certain arc, the same being measured on the scale *D*. This pendulum was a small, hollow glass bead, suspended by a silk fibre and trained delicately against the membrane. The bob carried a pointer for the scale *D*.

In the experiments the ball would be dropped against *A* some five or ten times, at intervals of about ten seconds, the corresponding deflections at *D* being noted and recorded; then the polarizer would be turned 90° and like observations noted. Again, 90° would be turned off, etc., etc.; occasionally the length of impulse arc would be changed, or more or less L-pieces applied, and in each case a large number of observations made.

In the experiments the initial pulse seemed to be followed by a series of vibrations in rapidly decreasing amplitudes; but it is believed that the initial pulse is equivalent to a genuine sound wave, or an undulation. Evidence of soundness of this view is found in the fact that the velocity of sound can be satisfactorily determined by similar pulses sent through tubes of 25 or 50 feet length. It was evident that the initial pulse only, was concerned in the first swing of pointer at *D*.

Observations were made with as few as one L in the polarizer, and two in the analyzer. But the results were small, averaging about 4 per cent. But during the noting of 455 individual results of observation the number of Ls in the polarizer ranged from 1 to 3 and in the analyzer 2 to 3. The average of all of these 455 observations gave a percentage of 8.87 hundredths of quenching of the polarized beam. That is to say, where the analyzer, in light, entirely quenches the polarized beam in turning through 90° , in the above 455 experiments only 8.87 hundredths of the polarized ray was quenched.

But as a higher percentage was looked for, the instrument itself was now examined for possible faults. The membranes were all found under considerable tension, whereas, of course, they should be perfectly free from it. After completely slackening, then, as was supposed, the experiments were continued with the following results:

POLARIZATION CONTINUED.

Coal Gas in L-tubes and Air outside.

Individual results.

Polarizer having 4 Ls and Analyzer 5 Ls.

=	+	=	+	=	+	=	+	=
6.0	6.0	8.0	5.0	6.5	5.8	6.2	6.0	6.0
6.0	6.0	7.5	5.5	6.5	5.5	6.0	5.2	6.2
6.2	6.0	7.0	5.3	6.3	5.6	6.5	5.2	7.2
6.1	5.8	7.5	5.2	6.5	5.2	6.4	5.1	6.2
6.0	5.7	7.0	5.0	6.2	5.7	6.3	5.5	6.1
6.1	5.5	7.2	5.2	6.6	6.0	6.3	5.1	6.1
6.2	5.8	6.5	5.0	6.8	5.2	6.2	5.2	6.1
6.0		7.0	4.9	6.2	5.2	6.5	5.4	6.5
		7.0	4.0	6.3	5.3	6.3	5.8	6.8
						6.3	5.2	6.4
Means	6.07	5.83	7.19	5.01	6.43	5.50	6.30	5.37

In this series each value given in any column is the number of divisions on the scale *D*, Fig. 3, of the deflection of the indicator pendulum bob. After obtaining one column of results the polarizer, *AB*, was turned 90° , and

the next column obtained. Thus the several columns were obtained. The = and + signify that the polarizer and analyzer are in parallel or perpendicular planes respectively.

On completing this series of observations air was passed into the L tubes, completely displacing the coal gas, so that the membranes were now suspended in mid air. Other conditions remained the same. The indicator pendulum now responded to the same impulses so slightly as to be barely observable, but not measurable, and they were apparently the same for the polarizer in all positions; that is to say, when air was upon both sides of the reflecting surfaces there was almost no appreciable reflection. This evidently should be the case, as the light membrane itself is now the chief cause of reflection. This, compared with the results obtained with gas in the tubes, shows that a considerable reflection is due to surfaces of separation of gases differing in density.

As regards the polarization, we observe that every mean under "—" is larger than any mean under "+," a fact which cannot be assumed accidental, nor explained on any other ground than polarization. To find the percentage of this in a single result, we have

Means of the above series.

$$\begin{array}{r} = \\ 6.07 \\ 7.19 \\ 6.43 \\ 6.30 \\ 6.34 \\ \text{Means, } 6.47 \\ 6.47 - 5.43 \\ \text{Per cent.} = \frac{6.47}{5.43} = 16.1. \end{array} \quad \begin{array}{r} + \\ 5.83 \\ 5.01 \\ 5.50 \\ 5.37 \\ 5.43 \\ 5.43 \end{array}$$

Whole number of observations, 80.

At this point the membranes were again examined and found to have appreciable tension, though supposed to have been entirely slackened at the beginning of the series of observations just cited. It seemed that the dampness in the gum used, expanded the membranes so that they became tight after thoroughly drying as the experiments went on. It was thereupon determined to slacken them with the utmost possible care, and continue the observations. The following table of individual results was then obtained in the manner explained above, the membranes being constantly watched for entire slackness.

POLARIZATION CONTINUED.

Coal Gas in tubes and Air outside of tubes.

Polarizer having 4 Ls and Analyzer 5 Ls.

Individual results.

=	+	=	+	=	+	=	+	=	+	=	+	=	+	=	+	=	+	=	+
1.2	0.9	2.0	1.0	1.5	1.0	1.6	1.0	1.8	1.4	1.5	0.9	2.0	1.0	1.5	1.0	2.1	1.2	1.5	1.0
1.5	1.0	2.1	1.2	1.5	1.0	1.4	1.1	1.5	1.2	1.5	1.0	2.1	1.2	1.5	1.0	1.8	1.1	1.5	1.0
1.5	1.0	1.8	1.1	1.8	1.0	1.8	1.1	1.5	1.4	1.5	1.0	1.8	1.1	1.5	1.0	2.0	1.0	1.5	1.0
1.5	1.0	2.0	1.0	1.3	1.0	1.9	1.2	1.7	1.2	1.5	1.0	2.0	1.0	1.5	1.0	2.0	1.0	1.5	1.0
1.5	1.1	2.0	1.0	1.3	1.0	1.9	1.0	1.7	1.1	1.5	1.1	2.0	1.0	1.5	1.0	2.0	1.0	1.5	1.0
		1.0	2.0	0.9	1.2	1.0	1.8	1.0	1.7	1.2	1.5	1.0	2.0	0.9					
			2.1	0.9	1.3	1.0	1.6	1.0	1.8	1.0									
					1.5	1.0			1.1	1.7	1.0								
									1.1	1.7									
									1.1	1.6									

Means 1.44 1.00 2.00 1.01 1.45 1.00 1.71 1.07 1.67 1.19 1.50 1.00 2.00 1.01

Polarizer turned 90° immediately following each column of results.

For gas displaced by air, other conditions the same; deflection, 0.00.

The smallness of these results, compared with those of the previous series, may be explained on the ground of extreme and entire slackness of the membranes; also the slackness is still further evinced by the fact that, when air displaced the gas no deflection of the indicator pendulum was observable in response to the impulses. Tense membranes would have turned the sound waves somewhat, and in the manner of the rebound of a drumstick from its drumhead.

To obtain the percentage of polarization effect, we have:

Mean of the above series.

=	+
1.44	1.00
2.00	1.01
1.45	1.00
1.71	1.07
1.67	1.19
1.50	1.00
2.00	1.01
Means, 1.68	1.04
Per cent.	1.68 — 1.04 = 38.1.
	1.68

Thus with the apparatus working most perfectly, the analyzer succeeded in extinguishing or quenching 38 per cent. of the polarized wave, a percentage too great to be mistaken. Considering the analyzer and polarizer as equally efficient, the real percentage of polarization by polarizer would be 62 per cent., and of the analyzer 62 per cent., as is evident from the fact that 62 per cent. of 62 per cent. is 38 per cent. Thus, either part of the apparatus obliterates over half of the wave attempting transmission, a fraction which would be unmistakably visible in polarized light.

These results establish the following facts for sound waves or for undulations, viz.:

1st. A decided reflection occurs at a surface separating two gases of different density, confirming the views of Henry and Tyndall in this regard.

2d. In repeated reflection from such surfaces the intensity of the final component varies with the relative positions of those surfaces, the same following the laws of polarization in light, from which we conclude that longitudinal undulations can be polarized.

With sound polarized, we complete the list of effects for longitudinal undulations which are known to light, viz.: radiation, shadows, reflection, refraction, diffusion, diffraction, interference and polarization; for the laws are common for like conditions, viz.: for intensity of radiation

in ambient space, $\frac{I}{d^2}$; in parallel space, $\frac{I}{d^1}$; in prismatic

space, like a tube, $\frac{I}{d^0}$; for shadows, reflection, refraction

and interference as well known; for diffusion, as when a steam whistle is sounded, filling the air with its resounding ring; for diffraction, as sound waves diverging rapidly after passing a narrow space between buildings, like light in passing a narrow slit and diverging; and, finally, for polarization, as above. In studying these comparisons we should recollect the vast difference between the properties of undulations in heavy, and ethereal media. Thus the wave length is very great and the velocity of propagation very small in sound as compared with light. This seems sufficient to account for the greater definition of shadows in light; but when a slit or an obstacle is made as narrow for light as for sound, in comparison to wave length, the diffraction divergence is probably about alike; that is, the divergence at a linear slit in light, or between two buildings in sound; or again the shadow of a silk fibre in light and a sound shadow of Bunker Hill monument, for instance, are about alike considering wave length. With these considerations it may be reasonable to expect incomplete or only partial polarization with such apparatus as employed above.

The conclusions to which we are conducted by the foregoing may be summed up as follows:

1st. That vibrations in extended media, produced from the action of a remote single centre of disturbance, can only be longitudinal, even in light.

2d. That vibrations will be to some extent transversal when due to two or more centres of disturbance not in the same line, as when two or more independent co-existent systems of undulations combine into one, or when a simple system is modified by such lateral disturbance as a reflection or a refraction.

3d. That undulations, to be in a condition called polarized, probably consist of vibrations which are transversal, and that no necessity exists for assuming vibrations transversal in front of a polarizer.

NOTE.—As regards longitudinal, oblique, transversal, etc., in the foregoing, the estimate is to be taken by comparing the direction of the line of vibration of a particle with that of propagation of the wave.

My acknowledgements are due to Mr. Clarence H. Wright, who, while a student in my physical laboratory last Spring, rendered valuable aid in the experimental work.

ASTRONOMY.

COMET (α) 1881—SWIFT.

The question of the best method of transmitting telegraphic announcements of astronomical discoveries has just been discussed by the leading European societies, and a system has been devised by which this information may be comprised in a message of sixteen words. Thinking that perhaps a better way existed of doing the same work, the Boston Scientific Society has adapted a telegraphic code to the needs of the occasion, and this system has just received a practical test. The announcements lie within the province of the Smithsonian Institution, and it was accordingly decided to transmit by cable the elements and ephemeris. These here given were computed at Dun Echt Observatory, in Scotland, by Drs. Copeland and Lohse, and have been distributed in this country to astronomers by special circulars of the Boston Scientific Society. That set which was computed at Boston, for the Society, by Mr. S. C. Chandler, Jr., has already been cabled to Europe, and distributed by mail, from the Observatory of Lord Crawford, to astronomers in England and the Continent.

The cablegram received at Boston consisted of sixteen words, and the translation is here appended. According to the same code, the announcement of discovery could be comprised in a message of seven words, which would itself contain check words against possible error in transmission.

The elements and ephemeris computed at Dun Echt, on Monday, May 9, were transmitted by cable to Boston in the following message: "Decimosexto erective contextual bewitchery anticly demonstrative courageously sputter arithmancy stomachical auriferous suety bayou synecdochically bissextile eminently." The translation of this message is as follows, viz:

ELEMENTS OF SWIFT'S COMET, 1881 (α).

Per. Passage, 1881, May 20.67, Greenwich Mean Time.

Long. Perihelion,	300	2	} Eq. 1881.0.
Long. Node,	124	54	
$\omega = \pi - \Omega$,	175	8	
Inclination,	78	48	
Log. $q = 9.7674$.	$q = 5854$.		
Motion direct.			

EPHEMERIS.

Greenwich, midnight.	(A.R.) h. m. s.	(DECL.) ° ' "	Brightness.
May 10,	0 38 32	+26 46	1.69
14,	56 48	21 35	
18,	1 17 32	15 54	
22,	40 48	9 55	2.32

Computed by Dr. R. Copeland and J. G. Lohse, from observations made at Dun Echt Observatory. The light at discovery is taken as unity.

By means of control-words in the message, it is absolutely known that the elements are those computed yesterday in Scotland, and it is proposed to cable in the same way the first elements and ephemerides of future comets, obtainable at either terminus, until the code has been most thoroughly tested.

J. RITCHIE, JR.

ON A METHOD OF ISOLATING THE MAMMALIAN HEART.

BY H. NEWELL MARTIN (Johns Hopkins University.)

To obtain a mammalian heart isolated from the rest of the body and keep it alive for a time sufficient to allow the examination of the effect of various conditions upon its activity has long been a physiological desideratum. The frog's heart has for years been the subject of minute study but hitherto the mammalian heart has been a baffling object. It seems to have been forgotten that while the frog's heart is a spongy structure having no arteries of its own, the mammalian heart is a dense organ dependent for its life on a continuous blood flow in its capillaries; and all attempts hitherto made, so far as I know, have been efforts to apply to the mammal the methods found successful with the frog, with merely the addition of arrangements adapted to keep up the comparatively high temperature at which the mammalian heart normally beats. By working in another way I have recently succeeded in keeping the mammalian heart alive for more than an hour, and beating with perfect rhythm and normal force; the organ is thus made almost as available for study as the heart of the frog. The method adopted is as follows: The animal having been narcotised and the chest opened, the aorta is tied just beyond its arch; then the trunk which, in the cat, gives origin to the right subclavian and the two common carotids, is ligatured close to its origin, and a cannula put in the left subclavian; finally, the inferior and superior venæ cavæ and the azygos vein, and the root of one lung are tied.

Artificial respiration is of course started so soon as the thorax is opened, and kept up henceforth. The course of the circulation is thus:—left auricle, left ventricle, commencement of aorta (and along the left subclavian to the cannula which is connected with a manometer), coronary system, right auricle, right ventricle, pulmonary vessels of one lung, and then back to the left auricle; in other words, the only section of the systemic circulation left is that through the vessels of the heart itself. Since the physiological actions taking place in the lung are among the best known of all occurring in the body, they may be eliminated, and we have practically an isolated and well-working living mammalian heart for study. The nerves going to the heart may be divided if desired, but that is hardly necessary as the want of blood-flow in the nerve centres of the body incapacitates them after a very short time, and they no longer are capable of exerting any influence on the heart. It is possible, however, that changes in the lung vessels may affect the results of experiments made on the heart's work under different conditions (*e. g.* when defibrinated blood is sent into it from a vein under various pressures, or when drugs are administered to it), and an investigation of the nerves, if any, governing the lung vessels must be undertaken as a preliminary to a further study of the direct action of various conditions on the heart's work.

EDINBURGH ROYAL SOCIETY.

There was a very large attendance at the meeting of the Edinburgh Royal Society held recently, to hear Professor Helmholtz, who was announced to make a communication "On Electrolytic Convection." Professor Helmholtz stated to the Society the results of certain later experiments which he had made in working out his theory of electrolytic convection—experiments which, he said, had succeeded better than his former ones had done. He had entered upon his experiments on account of certain objections that had been made to Faraday's electrolytic law, in connection with experiments which showed that a very feeble galvanic current could be kept up by moderate electro-motive force between platinum plates dipped in a slightly acidulated solution, even if the electro-motive force of the battery was not sufficient to decompose the water. He had found, in his earlier experiments, that the

only effect of the current was to absorb oxygen given off from the atmosphere and to form a new portion of water, and that the whole electrolytic effect was, not to decompose water and to produce a new quantity of elements—hydrogen and oxygen—but only that on the surface of one plate of platinum oxygen was collected and taken away from the surface of the other. He had, therefore, called these currents electrolytic convection. It was not really a decomposition, but only a transport of one of the products of electrolytic decomposition from one place to another. In his later experiments he had got rid of the atmospheric oxygen, by inclosing the whole electrolytic fluid in a sealed glass vessel. As a result, he had found that the smallest electromotive force, down to the thousandth of a Daniell's cell, produced a strong deflection which went immediately back to zero, that there was no continuous current, and that, if they broke the current, they got a deflection of the same character in the opposite direction to the first and direct one. Sir William Thomson, in speaking on the paper, stated that Professor Helmholtz's theory of electrolytic convection formed quite an era in electrolytic chemistry. Sir William himself made a communication "On the average pressure due to impulse of vortex-rings on a solid," following up inquiries suggested by Helmholtz's vortex theory. Professor Tait stated the result of certain calculations which he had made "On the crushing of glass by pressure," and which had been suggested by his recent inquiry as to error-corrections in the use of the *Challenger* thermometers. The result was that with cylindrical tubes made from ordinary Leith plate-glass, he found the glass gave way under a shear of about 1-250th, and that the strength of the tube was greater as the walls were thicker and the internal diameter decreased. But, however thick the walls might be, or however small the internal diameter, a pressure of 22 or 23 tons on the square inch would inevitably crush the strongest glass.

NOTES.

M. DAUDIGNY, electrical engineer in Paris, has sent to the Municipal Council a petition asking for authority to establish on the top of the Colonne de Juillet a large electric lamp fed by a magneto-electric machine of fifty horse-power. This enormous light is to be diffused by a large reflector of special construction.—*Nature*.

SIGNOR MANET, we learn from an Italian journal, whitens the albumen of blood by means of the electric light; which is projected by a system of mirrors and lenses, giving a strongly luminous effect. The time required varies according as the albumen is more or less separated from the fibrine. In general, 24 hours suffices to give complete decoloration.

PROFESSOR LOOMIS appears still to be experimenting in aerial telegraphy—telegraphing without wires—and it is now said that he proposes to establish communication, through the current which he claims is always found at a great altitude, between one of the highest peaks of the Alps, in Switzerland, and a similarly situated station on the Rocky Mountains, on this continent.

It has been found by M. Laurent that any ordinary good silvered glass mirror, plane, concave, or convex, and of any thickness, may be rendered a magical one by means of heat. A simple way of doing this is to heat a brass tube, and apply the end of it to the silvered face. If the mirror surface is opposite a screen, the section of the tube is reproduced in white; if the former is turned away from the screen, the image (which is seen only after removing the tube) is dark. A cold tube may be used with a hot mirror, and the experiment may be otherwise varied.

M. DUCHEMIN, the inventor of the compass with a circular magnet, now adopted in the French navy, has recently devised, for correcting compasses, a system of magnetic compensators. In place of the straight magnetic bars generally employed he uses magnets of an annular form. If we magnetise a steel ring, it may have two poles at opposite extremities of a given diameter and two neutral lines. Such rings—round, oval, or of any other form, and with or without interruption of continuity—may be utilised for the correction of a compass, by being placed either on the bridge of the vessel or in the binnacle.—*Revue Industrielle*.

Electrical Batteries for Medical Purposes.

THE WESTERN ELECTRIC MANUFACTURING CO.

CHICAGO, ————— NEW YORK,

MANUFACTURERS OF AND DEALERS IN

STATIONARY GALVANIC BATTERIES, PORTABLE HIGH ELECTROMOTIVE FORCE BATTERIES, CHLORIDE OF SILVER BATTERIES, GALVANO-CAUTERY BATTERIES, FARADAIC BATTERIES, POCKET BATTERIES, BATH APPARATUS, AND ELECTRODES.

Dr. Butler's Plumbago Rheostat,
PLIABLE SPONGE ELECTRODES, &c., &c.

CATALOGUES ON APPLICATION.

Address all Orders to

Western Electric Man'g Co.

38 Union Square,
— NEW YORK —
CITY.

It is successfully used by Schools, Colleges, Banks, Commission Houses, Mercantile Firms, Lawyers, Ministers, Copyists, and every person or firm desiring a number of duplicates of any document.

10,000 Pens now in Use and the Demand Steadily Increasing!

PRICE: No. 1, \$25.00; No. 2, \$30.00; No. 3, \$35.00.

Full Description and Samples of Work sent on Application.

EDISON'S ELECTRIC PEN AND PRESS,

38 UNION SQUARE, NEW YORK CITY.




EDISON'S Electric Pen & Duplicating Press,

An Extremely Useful Invention

FOR

DUPLICATING WRITTEN OR PRINTED MATTER.

8,000 Copies from a Single Writing, at the Rate of 400 an Hour.



**JAMES W. QUEEN & CO.
OPTICIANS,**

924 Chestnut St., Philadelphia, Pa.

The New \$50 Microscope
(See Illustration.)

Write for Catalogue of MAGNIFYING GLASSES, MICROSCOPES, MICROSCOPIC OBJECTS, MICROSCOPIC BOOKS, MICROSCOPIC OBJECT GLASSES, MICROSCOPIC ACCESSORIES, MOUNTING MATERIALS, &c., and mention this Journal.

Catalogues in 4 parts as follows:
Part 1st, Mathematical Instruments.
" 2d Optical Instruments.
" 3d Magic Lanterns and Slides.
" 4th Physical Instruments.

From our own experience we can confidently recommend to all Nervous and Debilitated persons

**Dr. CROSBY'S BRAIN AND NERVE FOOD,
THE VITALIZED PHOSPHITES,**
BECAUSE

It is largely prescribed by Physicians.

It is much taken by Clergymen, School Teachers, Lawyers, Students, and all who over-use the brain.

It cures Nervous Disorders, especially where it exists with Debility By curing Debility, it prevents Consumption.

Many who have been great sufferers with Nervousness, Sleeplessness and Neuralgia, speak of it in the highest praise.

It gives vigor to the mental and bodily growth of children, and makes a less fretful and happier childhood.

Every Wholesale Druggist in the U. S. buys it in large quantities.

SCIENCE

A WEEKLY JOURNAL OF SCIENTIFIC PROGRESS

Vol. II. No. 46, — — — — — May 14, 1881.

CONTENTS.

The "Zero Motor," (Edit.); Discovery of Fossil Forms in Meteorites, (Edit.); The Venom of Human Saliva; Scientific Societies of Washington; The Ultra-Gaseous or Radiant State of Matter, by Prof. H. S. Carhart; The Great Astronomical Telescope for the Vienna Observatory; Professor Draper's Photographs of the Nebula in Orion; The Polarization of Sound, by Prof. S. W. Robinson; On a Method of Isolating the Mammalian Heart, by H. Newton Martin; Edinburgh Royal Society; Notes, Astronomical, Chemical, Physical, &c., &c., &c.

SUBSCRIPTION, FOUR DOLLARS A YEAR.

SINGLE NUMBERS, TEN CENTS.

BRAIN AND NERVE FOOD.

VITALIZED PHOS-PHITES,

Composed of the Vital or Nerve-giving Principles of the Ox-Brain and Wheat-Germ.

IT RESTORES THE ENERGY LOST BY NERVOUSNESS OR INDIGESTION; RELIEVES LASSITUDE, ERRATIC PAINS AND NEURALGIA; REFRESHES THE NERVES TIRED BY WORRY, EXCITEMENT, OR EXCESSIVE BRAIN FATIGUE; STRENGTHENS A FAILING MEMORY, AND GIVES RENEWED VIGOR IN ALL DISEASES OF NERVOUS EXHAUSTION OR DEBILITY IT IS THE ONLY **PREVENTIVE** OF CONSUMPTION.

IT GIVES VITALITY TO THE INSUFFICIENT BODILY OR MENTAL GROWTH OF CHILDREN, PREVENTS FRETFULNESS AND GIVES QUIET, REST AND SLEEP. IT GIVES A BETTER DISPOSITION TO INFANTS AND CHILDREN, AS IT PROMOTES GOOD HEALTH TO BRAIN AND BODY.

PHYSICIANS HAVE PRESCRIBED 300,000 PACKAGES. For Sale by Druggists or by Mail, \$1.

F. CROSBY, 664 & 666 Sixth Ave., N. Y.—London, 137 A Strand.

DRAWING INSTRUMENTS

And Materials of all kinds.

OPTICAL INSTRUMENTS,

Microscopes, Telescopes, &c.

**G. S. WOOLMAN, 116 Fulton Street,
NEW YORK.**

Send for Illus. Catalogue, and mention this Journal.

Successful Novels.

MR. PERKINS' DAUGHTER.

By the MARCHIONESS CLARA LANZA.

Note.—"It may be well to mention, that the state of 'double consciousness' or 'Periodical Amnesia,' described in one of the principal personages of this story, is not by any means an isolated instance. It is, on the contrary, a phenomenon well known to neurological physicians."

**G. P. PUTNAM'S SONS,
27 West 23rd St., New York.**

CELLULOID EYE-GLASSES.

Representing the choicest selected Tortoise-Shell and Amber. The **LIGHTEST, HANDSOMEST AND STRONGEST** known. Sold by Opticians and Jewellers. Made by **SPENCER OPT. MFG CO.,
13 MAIDEN LANE, NEW YORK.**

American Journal of Science.

FOUNDED BY PROF. SILLIMAN IN 1818.

Devoted to Chemistry, Physics, Geology, Physical Geography, Mineralogy, Natural History, Astronomy, and Meteorology.

Editors: JAS. D. DANA, EDW. S. DANA and B. SILLIMAN. *Associate Editors:* Professors ASA GRAY, J. P. COOKE, JR., and JOHN TROWBRIDGE, of Cambridge; H. A. NEWTON and A. E. VERRILL, of Yale; and G. F. BARKER, of the University of Pennsylvania, Phila. Subscription Price \$6; 50 Cents a Number. Address the Proprietors.

JAMES D. & E. S. DANA, New Haven, Conn.

New York & New England R. R.

THROUGH TRAINS Via N.Y., N. H. & H. R. R. Between Boston and New York.

FROM GRAND CENTRAL DEPOT.

Leave New York, 11.00 A.M. Connects at Hartford.

" " " 4.00 P.M., with Elegant Through Car.

" " " 11.35 P.M., week days, with Through New Pullman Sleepers.

" " " 10.30 P.M., Sundays.

FROM DEPOT, FOOT OF SUMMER ST.

Leave Boston, 9.00 A.M., week days, with Through Cars to Grand Central Depot.

" " " 10.00 P.M., daily, with elegant new Pullman Sleepers.

G. P. PUTNAM'S SONS,

27 & 29 W. 23d St., N. Y.

Publish THE ELEMENTARY SCIENCE SERIES, 29 Vols., ready; THE ADVANCED SCIENCE SERIES, 14 Vols., ready; THE HOUSEHOLD SERIES OF ATLASES, from 75 cts. to \$25, and various other standard scientific books and works of reference.

Full catalogues sent on application. Orders filled on favorable terms for all American and Foreign Publications.

TAKE THE BEST.

THE INTERNATIONAL REVIEW.

NEW MONTHLY SERIES.

JOHN T. MORSE, JR., HENRY CABOT LODGE, Editors.

Price, 50 Cents a month; \$5.00 a year. Specimen copies sent post paid on receipt of 15 cents.

AGENTS WANTED who understand the character, scope and value of the REVIEW, to solicit subscriptions.

A. S. BARNES & CO.,

111 & 113 William Street, New York.

H. M. RAYNOR,

No. 25 Bond St.,

NEW YORK.

(Established
1859).



FOR ALL

Laboratory

AND

Manufacturing Purposes.

Wholesale and Retail.

Scrap Purchased.

Circulars by Mail.

BEST MEDIUM FOR ADVERTISERS.

SCIENCE

Circulates in EVERY STATE IN THE UNION, also Canada, Australia, Japan and Europe.

ALL UNIVERSITIES and COLLEGES.

Subscribers to "SCIENCE" are chiefly among those having few facilities for purchasing goods, and are buyers of Fancy Articles, Household Necessities, Jewelry, Musical Instruments, Dry Goods, Books, Prints, Scientific Instruments and Apparatus; Pure Chemical Substances, Mineral Waters, Novelties and Specialties of every kind.


Terms for Advertising moderate. Small advertisements on this page \$50 a year (52 insertions).

MEYROWITZ BROTHERS,

OPTICIANS,

297 Fourth Ave., N. Y.,

S. E. COR. 23D ST



Trial Cases, Ophthalmoscopes, Artificial Eyes, Clinical Thermometers, Microscopes, Electric Batteries, etc.
Special attention given to prescriptions of Oculists.

N. B.—An Illustrated Catalogue will be sent free upon request.

OPTICAL INSTRUMENTS.

AROMETERS, THERMOMETERS,
RAIN GAUGES, ETC., ETC.

**BENJ. PIKE'S SONS & CO.,
OPTICIANS,**

No. 972 Broadway, New York.

Illustrated Catalogue sent free on mentioning SCIENCE."

**R. & J. BECK,
MANUFACTURING OPTICIANS,
1016 Chestnut St., Philadelphia.**

MICROSCOPES,

And all Accessories of the HIGHEST GRADES at the LOWEST PRICES.
Illustrated Condensed Price List, 32 pages. Free.
Full Catalogue of 176 Pages. 15 cts.
Mention this Journal.

American Naturalist.

Devoted to the Natural Sciences in their widest sense.

PUBLISHED MONTHLY.

Terms, \$4.00 a year. Single numbers, 35 cents.

EDITORS:

Professor A. S. PACARD, JR., and Professor
E. D. COPE.

McCALLA & STAVELY, Publishers, Nos. 237-9
Dock Street, Philadelphia, Pa.